

On the other hand, applying  $\frac{\partial}{\partial q^i}$  to Hamilton-Jacobi gives the following "derived H-J equation:

$$(2) \quad 0 = \frac{\partial^2 S}{\partial t \partial q^i} + \frac{\partial H}{\partial q^i}(t, dS_t) + \sum \frac{\partial H}{\partial p_j} \frac{\partial^2 S}{\partial q^j \partial q^i}$$

Now  $\frac{\partial}{\partial q^j} \frac{\partial S}{\partial q^i} = \frac{\partial^2 S}{\partial q^i \partial q^j}$ . Hence subtracting the last two equations gives


(with suitable arguments)

$$(3) \quad \frac{\partial p_{i,c}}{dt} = \frac{\partial H}{\partial q^i}$$

This is the second half of Hamilton's equations.

In the last equation everything is to be regarded as a function of  $t$ .

The "suitable arguments" required to make this the case are indicated without ambiguity by the diagram of the functions involved.



It should be possible to make a systematic use of such mapping diagrams to indicate which (composite) arguments we intended in equations (such as the Hamilton and Hamilton-Jacobi equations).

Now consider the converse part of the theorem. Take a solution  $b$  of the first Hamilton equations; then equations (1) above hold. By hypothesis (1) implies (3); subtracting, (2) holds along  $b$ . But by the existence theorem for ordinary differential equations, there is a solution  $b$  through

each point of  $I \times C$ ; hence (2) holds at any point  $(t, q^1, \dots, q^n)$ . But

(2) states that 
$$\frac{\partial}{\partial q^i} (HJ(S)) = 0, \quad i = 1, \dots, n,$$

where  $HJ(S)$  denotes the left-hand side of the Hamilton-Jacobi equations.

*ie HJ(S) depends on t, so*

Therefore, there is a smooth function  $\theta: \mathbb{R} \rightarrow \mathbb{R}$  with

$HJ(S) = \theta \circ t: I \times C \rightarrow \mathbb{R}$ . Take a function  $\psi$  such that  $\frac{d\psi}{dt} = \theta$ . Then it is

not hard to see that

$$HJ(S - \psi) = 0.$$

This gives the conclusion of the theorem.

#### 44. Transformation to Equilibrium.

Now let  $Y$  be an  $n$ -dimensional manifold, and suppose

$S: \mathbb{R} \times C \times Y \rightarrow \mathbb{R}$  is a function such that everywhere

$$\det \left| \frac{\partial^2 S}{\partial q^i \partial y^j} \right| \neq 0$$

where  $q^i$  are coordinates for  $C$  and the  $y^j$  are coordinates for  $Y$ .

Let  $\mathbb{H}: \mathbb{R} \times C \times Y \rightarrow \mathbb{R} \times T^*C$  be given by

$$p_i^{\mathbb{H}} = \frac{\partial S}{\partial q^i}$$

$$t^{\mathbb{H}} = t$$

$$q^i{}^{\mathbb{H}} = q^i, \quad i = 1, \dots, n.$$

Thus the assumption on  $S$  is equivalent to saying that  $\mathbb{H}$  is regular

everywhere. Consider also the mapping  $\Psi: \mathbb{R} \times C \times Y \rightarrow \mathbb{R} \times T^*Y$

given by

$$\begin{aligned}
 t\Psi &= t \\
 y^i\Psi &= y^i \\
 x_i\Psi &= \frac{\partial S}{\partial y_i}, \quad i = 1, \dots, n,
 \end{aligned}$$

where the  $y^i$  are coordinates on  $Y$  and the  $x_i$  the corresponding (momentum) coordinates on  $T^*Y$ . This map is also regular everywhere.

Hence we have the diagram

$$\begin{array}{ccc}
 \mathbb{R} \times T^*C & \xleftarrow{\dots \chi \dots} & \mathbb{R} \times T^*Y \\
 \swarrow \textcircled{H} & & \nearrow \Psi \\
 \mathbb{R} \times C \times Y = N & & 
 \end{array}$$

and locally at least there is a map  $\chi$  from one time dependent phase space to another (compare §26).

Theorem. Take  $H: \mathbb{R} \times T^*C \rightarrow \mathbb{R}$ . For each point  $a \in Y$ , suppose that  $S: N \rightarrow \mathbb{R}$  satisfies the H-J partial differential equation for the function  $H$ . Let  $c: \mathbb{R} \rightarrow \mathbb{R} \times T^*Y$  be a curve of the form  $c(t) = (t, \text{const.})$ . Then the curve  $\chi c$  satisfies the Hamilton equations for  $H$  on  $\mathbb{R} \times T^*C$ .

Proof. It will be more convenient to look at everything in  $N$  rather than in  $\mathbb{R} \times T^*C$ . To do this we pull everything back locally by  $\textcircled{H}^{-1}$ . Thus we are interested in the curve  $\Psi^{-1}c$  and the function  $H\textcircled{H}$ .

Let  $x_i$  denote the coordinates as above. Then  $x_i = \frac{\partial S}{\partial y_i}$  on  $N$ .

Taking  $\frac{d}{dt}$  of this, we get

$$0 = \frac{\partial^2 S}{\partial y^i \partial t} + \sum \frac{\partial^2 S}{\partial y^i \partial q^j} \frac{dq^j}{dt} + \sum \frac{\partial^2 S}{\partial y^i \partial y^j} \frac{\partial y^j}{\partial t} .$$

The third sum vanishes because  $\frac{\partial y^i}{\partial t} = 0$ . Since we are assuming

$$\frac{\partial S}{\partial t} + H = 0$$

on  $N$  (more precisely, we should write  $H \circledast$  instead of  $H$ ), applying

$\frac{\partial}{\partial y^i}$  yields

$$\frac{\partial^2 S}{\partial t \partial y^i} + \sum \frac{\partial H}{\partial p^j} \frac{\partial^2 S}{\partial q^i \partial y^j} = 0,$$

which holds on the curve  $\Psi^{-1}c$ . Hence, again on the curve,

$$\sum \frac{\partial^2 S}{\partial q^i \partial y^j} \left( \frac{dq^j}{dt} - \frac{\partial H}{\partial p_j} \right) = 0.$$

Since the determinant of  $\left( \frac{\partial^2 S}{\partial q^i \partial y^j} \right)$  was assumed to be non-zero, this

means that

$$\frac{dq^j}{dt} = \frac{\partial H}{\partial p_j}$$

holds for all  $j$ . This is the first Hamilton equation. By the previous theorem, we get the remaining half of the Hamilton equations.

For fixed  $t_0$ , the submanifolds of  $N$  of the form  $t_0 \times C \times Y$  have a symplectic form given by

$$\sum \frac{\partial^2 S}{\partial q^i \partial y^j} dq^i \wedge dy^j.$$

By the theorem proved in § 26 of Part I, the functions  $\circledast$  and  $\Psi$  are symplectic mappings, whence the usual symplectic structures are taken on  $T^*C$  and  $T^*Y$ .

In the theorem just proved, the trajectories  $c$  in  $\mathbb{R} \times T^*Y$  are constant in  $T^*Y$ . Hence one says that the map  $\chi^{-1}$  of the theorem transforms the Hamiltonian  $H$  "to equilibrium".

45. Characteristics.

The previous results indicate a close relation between the Hamilton-Jacobi equation, a partial differential equation, and Hamilton's equations, a system of ordinary first order differential equations. This is a special case of the theory which relates a first order partial differential equation to its characteristics, which are solutions of a corresponding system of ordinary first order differential equations.

- Sources: a) Courant and Hilbert, Methods of Mathematical Physics, II  
 b) Caratheodory, Calculus of Variations and PDE's of 1<sup>st</sup> order, Part I.

The case of the arbitrary first order equation will be reached in stages. We first consider the linear case, involving the following functions on a configuration space  $C$ :

$$\mathbb{R} \xrightarrow[c]{q^1, \dots, q^n} C \xrightarrow[u]{q^1, \dots, q^n} \mathbb{R},$$

$$(1) \quad \sum_{i=1}^n a_i \frac{\partial u}{\partial q^i} = bu + d, \quad \begin{cases} a_i: C \rightarrow \mathbb{R}, \\ b: C \rightarrow \mathbb{R}, \\ d: C \rightarrow \mathbb{R}. \end{cases}$$

Equation (1) for the linear case has  $a_i, b, d$  functions of position in  $C$ .

The  $a_i$  determine a vector field  $X = \sum_{i=1}^n \frac{\partial}{\partial q^i}$  on  $C$  which appears in  $a_i$

the following coordinate independent form of (1):  $L_X u = bu + d$ . Call  $c$

a characteristic curve of the PDE (1) when


$$(2) \quad \frac{d(q^i c)}{dt} = a^i, \quad i = 1, \dots, n.$$

Thus the characteristics are the trajectories of the vector field  $X$ . In view of the definition of  $X$ , this equation can be written as

$$(2') \quad \frac{du}{dt} = bu + d.$$

First suppose that  $b = d = 0$ . Then if the function  $u$  satisfies the PDE (1),  $L_X u = 0$  it is constant along the characteristics of (1). More generally, for any  $b$  and  $d$ , the equation states that the values of  $u$  along a characteristic are determined by the value at any one (initial) point there. This suggests that we can obtain a solution  $u$  by taking initial values along a suitable set  $S$ , and then prolong these values by solving (2').

More explicitly, find a submanifold  $S$  of dimension  $n-1$  transverse to  $X$  (i.e., with  $T_a C = T_a S \oplus \mathbb{R}X(a)$  at each point  $a$  of  $S$ ). According



to the basic theorem on the integration of (smooth) vector fields, the trajectories of  $X$  through  $S$  cover some neighborhood of  $S$ , determining on some neighborhood of any  $\overset{\circ}{S}$  a unique function  $u$  which agrees on  $\overset{\circ}{S}$  with some chosen  $u_0 : \overset{\circ}{S} \rightarrow \mathbb{R}$ , and which satisfies (2) along characteristics (trajectories of  $X$ ) on the neighborhood. Here  $\overset{\circ}{S}$  is an open submanifold of  $S$  with compact closure in  $S$ . In local coordinates it is immediate that, for smooth  $u_0$ , the function  $u$  is smooth and satisfies (1). So we have found a local solution.

Next we consider a first order P.D.E. in an unknown  $u$ , of the form

$$(1) \quad \sum_{i=1}^n a_i(u, q^1, \dots, q^n) \frac{\partial u}{\partial q^i} = b(u, q^1, \dots, q^n), \quad u = u(q^1, \dots, q^n).$$

This is linear in all the partial derivatives of  $u$ , but not in  $u$  itself, hence is said to be quasilinear. We interpret the  $q^1, \dots, q^n$  as coordinates in an  $n$ -dimensional configuration space  $C$ , so that  $u: C \rightarrow \mathbb{R}$ . The equation thus has the form

$$(1') \quad \sum_{i=1}^n a_i \frac{\partial u}{\partial q^i} = b, \quad \mathbb{R} \times C \begin{array}{c} \xrightarrow{a_1} \\ \vdots \\ \xrightarrow{a_n} \\ \xrightarrow{b} \end{array} \mathbb{R},$$

for given coefficient functions  $a_i$  and  $b$ .

We plan to reduce this to the previous case for a linear P.D.E. in an unknown  $v: \mathbb{R} \times C \xrightarrow{\mathbb{R}} \mathbb{R}$  in one more variable, constructing the function  $u$  via its graph  $\hat{u}: C \xrightarrow{\hat{u}} \mathbb{R} \times C$ . Let  $r: \mathbb{R} \times C \rightarrow \mathbb{R}$  be the projection on the first coordinate. We introduce a function  $v: \mathbb{R} \times C \xrightarrow{v} \mathbb{R}$  defined by  $v = u - r$ . Now  $\frac{\partial v}{\partial r} = -1$  and  $\frac{\partial v}{\partial q^i} = \frac{\partial u}{\partial q^i}$  (while on hypersurfaces

$v = 0$  we will have  $a_i(u, q^i) = a_i(r, q)$  and  $b(u, q) = b(r, q)$ ) so that (1)

becomes

$$0 = \sum_{i=1}^n a_i \frac{\partial u}{\partial q^i} - b = \sum_{i=1}^n a_i \frac{\partial v}{\partial q^i} + b \frac{\partial v}{\partial r} = \left( \sum_{i=1}^n a_i \frac{\partial}{\partial q^i} + b \frac{\partial}{\partial r} \right) v.$$

This is a homogeneous linear P.D.E. in  $v: \mathbb{R} \times C \rightarrow \mathbb{R}$ . Its characteristics are thus given by a suitable vector field  $\hat{X}$ . Indeed we now define the vector field  $\hat{X}$  on  $\mathbb{R} \times C$  by  $\hat{X} = \sum_{i=1}^n a_i \frac{\partial}{\partial q^i} + b \frac{\partial}{\partial r}$ , then (on

the hypersurface  $v = 0$ ) the equation (1) becomes:

$$0 = \left( \sum_i a_i \frac{\partial}{\partial q^i} + b \frac{\partial}{\partial r} \right) v = \langle dv, \hat{X} \rangle .$$

So  $\hat{X}$  at each point is in the tangent plane to the hypersurface at that point, and the trajectories of  $\hat{X}$  remain in the hypersurface. Moreover:

Proposition. For  $x_0 \in \mathbb{R} \times C$ , let  $v: \mathbb{R} \times C \rightarrow \mathbb{R}$  with

$$\begin{cases} \langle dv, \hat{X} \rangle = 0 \\ \left. \frac{\partial v}{\partial r} \right|_{x_0} \neq 0, v(x_0) = 0. \end{cases}$$

Then the function  $u: N_{x_0} \rightarrow \mathbb{R}$  such that  $v(q^1, \dots, q^n, u) = 0$ , constructed for a suitable neighborhood  $N_{x_0} \subset C$  via the implicit function theorem, is a solution of the P. D. E. (1).

Proof. By the construction of  $u$ ,


$$\begin{aligned} 0 &= \frac{\partial}{\partial q^i} (v \circ \hat{u}) & C &\xrightarrow{\hat{u}} \mathbb{R} \times C \xrightarrow{v} \mathbb{R}, \\ &= \frac{\partial v}{\partial q^i} + \frac{\partial v}{\partial r} \frac{\partial u}{\partial q^i} \end{aligned} \text{ for each } i, \text{ so that we have}$$

$$0 = \langle dv, \hat{X} \rangle = \sum_{i=1}^n a_i \frac{\partial v}{\partial q^i} + b \frac{\partial v}{\partial r} = \sum_{i=1}^n -a_i \frac{\partial v}{\partial r} \frac{\partial u}{\partial q^i} + b \frac{\partial v}{\partial r} .$$

Therefore for  $\frac{\partial v}{\partial r} \neq 0$ ,  $u$  will be a solution of (1), q. e. d.

Let  $S_0$  be a submanifold of  $C$  of dimension  $n-1$ , and let  $u_0: S_0 \rightarrow \mathbb{R}$  be a smooth function. Through  $S_0$  in  $\mathbb{R} \times C$  pass the vertical hypersurface  $T = \{(r, x) \mid r \in \mathbb{R}, x \in S_0\}$ . Define  $v_0: T \rightarrow \mathbb{R}$  by  $v_0(r, x) = u_0(x) - r$ . Suppose that the characteristic field  $\hat{X}$  is transverse to  $T$  at a point  $x_0$  of the  $n-1$  dimensional submanifold  $S$  on which  $v_0 = 0$ . Then, it is immediate in local coordinates (see the

figure below) that the trajectories of  $\hat{X}$  through some neighborhood in  $S$  of  $x_0$  determine an  $n$ -dimensional submanifold  $\Sigma$  of  $\mathbb{R} \times C$  (locally unique). Moreover,  $\Sigma$  is the graph of the desired solution  $u$ . For the



(Note: in the figure,  $C$  is the horizontal plane and the  $\mathbb{R}$  axis is the vertical.)

fact that  $\hat{X}$  is transversal to  $T$  at  $x_0$  implies that the function  $v(x, t) = v_0(x)$  (where  $x \in T$  and  $\hat{X} = \frac{\partial}{\partial t}$ ) is well-defined on a neighborhood of  $x_0$  in  $\mathbb{R} \times C$  and satisfies the conditions of the previous proposition  $(\frac{\partial v}{\partial r} \Big|_{x_0} = -1 \neq 0)$ .

Any point of  $\mathbb{R} \times C$  at which  $\hat{X}$  is non-vertical lies on the graph of such solutions. Explicitly, the hypersurface  $S_0 \subset C$  may be described as the locus where some smooth function  $f: C \rightarrow \mathbb{R}$  is constant (i.e., as a level hypersurface of  $f$ ). Then the vertical hypersurface  $T$  is

$$T = \{(r, y) | f(y) = f(y_0)\}$$

for  $y_0$  a fixed and  $y$  any point of  $C$ .

We have proved

Theorem. For smooth functions  $a_1, \dots, a_n, b: \mathbb{R} \times C \rightarrow \mathbb{R}$  let  $S_0 \subset C$  be defined by a point  $y_0 \in C$  and a smooth function  $f: C \rightarrow \mathbb{R}$  as

$$S_0 = \{y \mid y \in C \text{ and } f(y) = f(y_0)\}.$$

If  $u_0: S_0 \rightarrow \mathbb{R}$  is a smooth function satisfying the "transversality"

condition

$$\sum a_i(u_0(y_0), y_0) \frac{\partial f}{\partial q^i} \neq 0,$$

then in some neighborhood of  $y_0$  there is a unique solution  $u$  of the

P.D.E. 
$$\sum a_i \frac{\partial u}{\partial q^i} = b$$
 with values  $u_0$  on  $S_0$ .

#### 46. The General First Order P.D.E.

Consider an equation

$$(1'') \quad E(u, q^1, \dots, q^n, \frac{\partial u}{\partial q^1}, \dots, \frac{\partial u}{\partial q^n}) = 0$$

in an unknown function  $u: C \rightarrow \mathbb{R}$ , where  $q^1, \dots, q^n$  are local coordinates

in the configuration space  $C$ . We can regard the "equation"  $E$  as a

given (smooth) function  $E: \mathbb{R} \times T^*C \rightarrow \mathbb{R}$ . The differential  $du$  is a func-

tion  $C \rightarrow T^*C$ ; we also have  $d'u: C \rightarrow \mathbb{R} \times T^*C$  given locally as

$$(q^1, \dots, q^n) \rightarrow (u, q^1, \dots, q^n, \frac{\partial u}{\partial q^1}, \dots, \frac{\partial u}{\partial q^n}).$$

Thus the equation (1'') becomes  $E \circ d'u = 0$ . If  $r: \mathbb{R} \times T^*C \rightarrow \mathbb{R}$  is the

projection on the first factor, then  $\frac{\partial}{\partial q^i}$  applied to (1'') yields the  $i^{\text{th}}$

derived P.D.E.

$$\frac{\partial E}{\partial r} \frac{\partial u}{\partial q^i} + \frac{\partial E}{\partial q^i} + \sum_{j=1}^n \frac{\partial E}{\partial p_j} \frac{\partial}{\partial q^i} \left( \frac{\partial u}{\partial q^j} \right) = 0, \quad i = 1, \dots, n.$$

Interchanging the order of partial derivatives, this is

$$(2) \quad \frac{\partial E}{\partial r} \frac{\partial u}{\partial q^i} + \frac{\partial E}{\partial q^i} + \sum_{j=1}^n \frac{\partial E}{\partial p_j} \frac{\partial}{\partial q^j} \left( \frac{\partial u}{\partial q^i} \right) = 0.$$

By way of motivation, observe that the  $i^{\text{th}}$  equation of (2) may now be regarded as a quasilinear P.D.E. in the unknown  $p_i = \frac{\partial u}{\partial q^i}$ . The

characteristics of this quasilinear equation are then given by the

$(n+1)$ -dimensional vector field (see above):

$$\hat{X} = \sum_j \frac{\partial E}{\partial p_j} \frac{\partial}{\partial q^j} + \left( - \frac{\partial E}{\partial q^i} - \frac{\partial E}{\partial r} p_i \right) \frac{\partial}{\partial r}.$$

The differential equations of these characteristics are then the  $n+1$  equations

$$\frac{dq^j}{dt} = \frac{\partial E}{\partial p_j}, \quad j = 1, \dots, n,$$

$$\frac{dp^i}{dt} = - \frac{\partial E}{\partial q^i} - \frac{\partial E}{\partial r} p_i.$$

As  $i$  varies, the first  $n$  equations are the same. Note also that this reduces to Hamilton's equations when  $\frac{\partial E}{\partial r} = 0$ .

Our actual interpretation of (2) will be slightly different, as an equation on  $\mathbb{R} \times T^*C$  itself, with characteristics in  $\mathbb{R} \times T^*C$  which are solution curves of:

$$(3) \quad \frac{dq^j}{dt} = \frac{\partial E}{\partial p_j}, \quad \frac{dp_i}{dt} = - \frac{\partial E}{\partial q^i} - \frac{\partial E}{\partial r} p_i,$$

$$\frac{dr}{dt} = \sum_j p_j \frac{\partial E}{\partial p_j}.$$

The third set of equations is included since  $E = E(r, q, p)$  is constant along

*Think of this as a Hamiltonian system on  $\mathbb{R} \times T^*C$ . The characteristic curves are the integral curves of the vector field  $\hat{X}$ .*

trajectories of the vector field

$$(3') \quad X_E = \sum \frac{\partial E}{\partial p_j} \frac{\partial}{\partial p_j} + \sum_j \left( -\frac{\partial E}{\partial q^j} - p_j \frac{\partial E}{\partial r} \right) \frac{\partial}{\partial p_j} + \sum p_j \frac{\partial E}{\partial p_j} \frac{\partial}{\partial r}$$

on  $T^*C \times \mathbb{R}$ . For


$$L_{X_E} E = \sum_j \frac{\partial E}{\partial p_j} \frac{\partial E}{\partial p_j} + \left( \sum_j -\frac{\partial E}{\partial q^j} \frac{\partial E}{\partial p_j} - \sum p_j \frac{\partial E}{\partial r} \frac{\partial E}{\partial p_j} \right) + \sum p_j \frac{\partial E}{\partial p_j} \frac{\partial E}{\partial r} = 0.$$

For our previous cases, the last equation disappears, while the middle equations all collapse to the equation for  $r$  (reabeled  $p_{1c}$ ) from the quasilinear case, giving trajectories "parallel" to those of the earlier cases. We state our existence theorem in the form:

Theorem: Given in  $C$  a compact submanifold  $S_0$  of dimension  $n-1$  and initial values  $u_0$  of  $u$  on  $S_0$  such that a certain determinant (which appears as (5) below) does not vanish, then there exists an open set  $U \supset S_0$  and a smooth function  $u: U \rightarrow \mathbb{R}$  which satisfies  $E$  on  $U$  and agrees on  $S_0$  with  $u_0$ .

It will be clear from the proof that the conditions on the initial surface could be taken as before, and that the determinant condition corresponds to our previous transversality condition, with no loss of applicability.

Proof. We operate in  $\mathbb{R} \times T^*C$ , where we already have defined in (3) the characteristic curves. In the submanifold  $T$  of dimension  $2n$  above  $S_0$ , we distinguish a surface  $S$  which will correspond to  $\hat{U} \cap T$ . This submanifold (diffeomorphic to  $S_0$ ) with local coordinates  $x^1, \dots, x^{n-1}$  embedded by  $v: S_0 \xrightarrow{v} S \subseteq T$ , should have



Proof. In the configuration space  $C$  we have local coordinates  $q^1, \dots, q^n$ , an initial manifold  $S_0 \subset C$  of dimension  $l$  and initial values  $u_0: S_0 \rightarrow \mathbb{R}$ . On  $\mathbb{R} \times T^*C$  we have  $2n+1$  local coordinates  $r, q^1, \dots, q^n, p_1, \dots, p_n$ . We can define a map  $v: S_0 \rightarrow \mathbb{R} \times T^*C$ ; this amounts to choosing "initial" values of  $r, q^i$  and  $p_j$  along  $S_0$ . Specifically we make  $r \circ v = u_0, q^i \circ v = q^i$  and then we choose  $p_1, \dots, p_n$  so that

$$E = 0, \quad dr - \sum p_i dq^i = 0$$

both along  $S_0$ . The last condition on  $dr$  may be written in terms of local parameters  $x^1, \dots, x^{n-1}$  on the  $(n-1)$ -manifold  $S_0$  as

$$(4) \quad 0 = du_0 - \sum_{i=1}^n p_i dq^i = \sum_k \left( \frac{\partial u_0}{\partial x^k} - \sum_i p_i \frac{\partial q^i}{\partial x^k} \right) dx^k.$$

Hence  $p_1, \dots, p_n$  are determined uniquely along  $S_0$  if

$$(5) \quad \begin{vmatrix} \frac{\partial E}{\partial p_1} & \frac{\partial E}{\partial p_2} & \cdots & \frac{\partial E}{\partial p_n} \\ \frac{\partial q^1}{\partial x^1} & \cdots & \frac{\partial q^n}{\partial x^1} \\ \vdots & & & \\ \frac{\partial q^1}{\partial x^{n-1}} & \cdots & \frac{\partial q^n}{\partial x^{n-1}} \end{vmatrix} \neq 0 .$$

*Handwritten scribble*

This condition may be readily satisfied, since we can assume that the first row is nowhere zero. (This amounts to assuming that the given partial differential equation effectively involves at least one of the partial derivatives  $p_i = \frac{\partial u}{\partial q^i}$ ). Given such a first row, the submanifold  $S_0$  can be chosen to make (5) hold; for example, if  $\frac{\partial E}{\partial p_n} \neq 0$  we can choose the submanifold  $S_0$  given locally by the equation  $q^n = 0$ , with local coordinates  $x^1 = q^1, \dots, x^{n-1} = q^{n-1}$ ; then the determinant (5) is simply  $(-1)^n \frac{\partial E}{\partial p_n}$ . Indeed, the condition (5) is then exactly the condition that  $S_0$  be transversal to the projection of  $X_E$ .

I?

We now have  $v: S_0 \rightarrow \mathbb{R} \times T^*C$ , with image an  $(n-1)$ -manifold  $S$  in  $\mathbb{R} \times T^*C$ ; moreover one can show  $S$  transversal to the characteristic vector field  $X_E$ . Therefore the trajectories of  $X_E$  through points of  $S$  fill up locally a manifold  $T$  of dimension  $n$ . Now  $E = 0$  holds along  $S$ , so by the properties earlier established for characteristics it holds along  $T$ . In other words,  $T$  gives the graph of functions  $u, q^1, \dots, q^n, p_1, \dots, p_n$  on  $C$  (or on a neighborhood of  $S_0$  in  $C$ ) which satisfy  $E(u, q^1, \dots, q^n, p_1, \dots, p_n) = 0$ .

What remains to be verified is that  $p_i = \frac{\partial u}{\partial q_i}$  for  $i = 1, \dots, n$  on this manifold  $T$ . If  $\nu: T \rightarrow \mathbb{R} \times T^*C$  is the inclusion map, this amounts to showing that the induced 1-form  $\theta = \nu^*(dr - \sum p_i dq^i)$  is zero on  $T$ . We may calculate  $\theta$  in local coordinates  $x^1, \dots, x^{n-1}$  (on  $S_0$ ) and  $t$  (the parameter along the trajectories of  $X_E$ ) as

$$0 = \sum_{k=1}^{n-1} \left( \frac{\partial r}{\partial x^k} - \sum p_i \frac{\partial q^i}{\partial x^k} \right) dx^k + \left( \frac{\partial r}{\partial t} - \sum p_i \frac{\partial q^i}{\partial t} \right) dt .$$

The last term is zero by the equation (3) for the characteristics. It thus remains to show that

$$D_k = \frac{\partial r}{\partial x^k} - \sum p_i \frac{\partial q^i}{\partial x^k} , \quad k = 1, \dots, n-1$$

is zero. But  $D_k = 0$  on  $S_0$  by the choice of the initial values of  $p_i$ , while a calculation with the equations (3) shows

$$\frac{\partial D_k}{\partial t} = \frac{\partial E}{\partial x^k} + \frac{\partial E}{\partial r} D_k = \frac{\partial E}{\partial r} D_k .$$

This is a linear first order differential equation for  $D_k$  as a function of the parameter  $t$ , with initial values zero on  $S_0$ . Hence (by the uniqueness of the solutions of such equations)  $D_k = 0$ , q. e. d..

47. Contact Manifolds. The use of the characteristic vector field  $X_E$  for the partial differential equation  $E$  raises the following question. For a symplectic manifold any two smooth functions  $f$  and  $g$  have a Poisson bracket given by

$$\{f, g\} = L_{X_f} g = -L_{X_g} f .$$

